

Markov

$$X \geq 0 \Rightarrow P(X \geq a) \leq \frac{EX}{a}$$

$$P(X=x) = \sum_{y \in Y(x)} P(X=x | Y=y) P(Y=y)$$

Fonction Rischier

$$h(x) = \frac{f(x)}{S(x)} \quad S(x) = \int_0^x h(t) dt$$

$$P(X \leq x + \Delta x | X > x) \approx \Delta x h(x)$$

Bayes

$P(E|H_2)$ association.

$$\frac{P(H_2|E)}{P(H_1)} = \max_i \frac{P(H_i|E)}{P(H_i)} \Leftrightarrow \text{la s'ila plus association.}$$

$$P(H_2|E) = \max_i P(H_i|E) \text{ He she divanta plus probab.}$$

$$\sum_{i=1}^m x^{i-1} = \frac{1-x^m}{1-x}$$

Cauchy

$$P(|X - EX| > k\sigma) \leq \frac{1}{k^2} \quad k \geq 1$$

$$P(|X| > c) \leq \frac{EX^2}{c^2} \quad c > 0$$

convolution

$$P(X+Y=z) = \sum_{y \in Y(z)} P(X=z-y) P(Y=y) \quad \text{Babou } f_{X+Y}^{(2)}(z) = \int_{-\infty}^{\infty} f_X(z-y) f_Y(y) dy$$

h continue $\Rightarrow$ intersection

- 1) $h(x) > 0$
- 2) $\int_0^{\infty} h(x) dx = +\infty$

Binomiale

$$P_k = \binom{n}{k} p^k (1-p)^{n-k} \quad K=0,1,\dots,n$$

K teste in n lanci

$$EX = np, \quad Var X = npq$$

$X = \# \text{ teste}$

Uniforme

$$U[a,b] \Rightarrow EX = \frac{a+b}{2}, \quad Var X = \frac{(b-a)^2}{12}$$

Poisson

$$P_k = e^{-\lambda} \frac{\lambda^k}{k!}, \quad \lambda > 0$$

K teste in $n = \infty$ lanci

$EX = \lambda$; ~~Var X = \lambda~~

$X+Y \sim \text{Pois}(\lambda_1 + \lambda_2)$; $P(X=x | X+Y=n) \sim \text{Binom}(n, \frac{\lambda_1}{\lambda_1 + \lambda_2})$

$X, Y \text{ indep.}$

Geometrica

$$P_m = p(1-p)^{m-1}, \quad EX = \frac{1}{p}, \quad Var = \frac{q}{p^2}, \quad X = \# \text{ lanci}; P(X \geq m) = q^{m-1}$$

testa dopo m lanci; assenza memoria $P(X > x_0 + x | X > x_0) = P(X > x_0)$

Purel

$$P(X=n) = \binom{n-1}{k-1} p^k (1-p)^{n-k}$$

K teste dopo n lanci

$X_i = \# \text{ lanci per ottenere l'i-esimo testa}; X = \sum_{i=1}^K X_i$

Esponenziale

$$f(x) = \lambda e^{-\lambda x}, \quad x \geq 0$$

$f(x) = 0, \quad x < 0$

$$EX^2 = \frac{2}{\lambda^2}$$

Normale

$$f(x) = \frac{1}{\sqrt{2\pi} \sigma} e^{-\frac{1}{2} \left(\frac{x-\mu}{\sigma} \right)^2}$$

$$EX = \mu, \quad Var X = \sigma^2$$

$$\Phi(x) = 1 - \Phi(-x)$$

$$\Phi(x) - \Phi(-x) = 2\Phi(x) - 1$$

$$\Phi(0) = \frac{1}{2}$$

$$Y = ax + b$$

$$Var Y = a^2 Var X = a^2 \sigma^2$$

$$P(|X - \mu| \leq k\sigma) = 2\Phi(k) - 1$$

Hypergeom

$$P_k = \frac{\binom{m}{k} \binom{n-m}{n-k}}{\binom{n}{n}}$$

surv substitution

$$X = E_1 + \dots + E_m; \quad E_i \text{ sono scambiabili}$$

Multinomiale

$$\{E^{(1)}, \dots, E^{(m)}\}$$

$$P(X_1=x_1, \dots, X_m=x_m) = \binom{n}{x_1, \dots, x_m} p_1^{x_1} \dots p_m^{x_m}$$

$x_1 + \dots + x_m = n$

$$EX = \frac{1}{\lambda}; \quad Var = \frac{1}{\lambda^2}$$

$$min(X, Y) \sim \text{Exp}(\lambda_1 + \lambda_2)$$

$X, Y \text{ indep.}$

$$Z = \max(X, Y) \Rightarrow P(Z \leq z) = P(X \leq z) P(Y \leq z)$$

$$Y = ax + b$$

$$Var Y = a^2 Var X = a^2 \sigma^2$$

$$P(|X - \mu| \leq k\sigma) = 2\Phi(k) - 1$$